

The climate response to CO_2 of the Hadley Centre coupled AOGCM with and without flux adjustment

by

J M Gregory and J F B Mitchell

CRTN 78

January 1997

CLIMATE
RESEARCH
TECHNICAL
NOTE

Hadley Centre
Meteorological Office
London Road
Bracknell
Berkshire RG12 2SY

The climate response to CO_2 of the Hadley Centre coupled AOGCM with and without flux adjustment

J. M. Gregory and J. F. B. Mitchell

27th January 1997

This paper has been submitted to *Geophysical Research Letters* and can be read on the MetWeb in PostScript at `http://hc0500/~hadsa/hadcm2d/hadcm2d.ps.gz`

Abstract

Coupled ocean-atmosphere general circulation models have a tendency to drift away from a realistic climatology. The modelled climate response to an increase of CO_2 concentration may be incorrect if the simulation of the current climate has significant errors, so in many models, including ours, the drift is counteracted by applying prescribed fluxes of heat and fresh water at the ocean–atmosphere interface in addition to the calculated surface exchanges. Since it does not have a good physical basis, the use of this technique of "flux adjustment" itself introduces some uncertainty in the simulated response to increased CO_2. We find that the global-average temperature response of our model to CO_2 increasing at 1% per year is about 30% less without flux adjustment than with flux adjustment, while the geographical patterns are similar. The reduced response is due to more effective vertical transport of heat into the ocean, and a somewhat smaller climate sensitivity, both of which relate to differences in the initial climates. The response of the model with flux adjustment is more credible because it has a more realistic initial climate. The geographical similarity indicates that flux adjustment is not causing any gross distortion of the response.

Introduction

Coupled atmosphere-ocean general circulation models (AOGCMs) are the most comprehensive tools available for time-dependent climate-change experiments. Although the component atmosphere and ocean GCMs may separately produce acceptable simulations when forced at the surface with externally supplied realistic boundary conditions, significant errors in the climatology typically develop when the two are coupled to make an AOGCM. This presents a problem: if the simulation of present conditions (the "control" climate) is unrealistic, the response of the simulated system to a perturbation may not be correct, because the climate system is non-linear. A simple illustration of this can be found in the global-mean radiative equilibrium $S = \sigma T^4$, where S is the radiative flux into the climate system, and T the surface temperature required for equilibrium, assuming black-body emission. Increasing CO_2 increases S, and $\Delta T/\Delta S = T/4S$. Hence

if T is wrong in the control, ΔT will also be wrong. As we shall see, there are other non-linearities which are more important.

Many AOGCMs (see *e.g.* Gates *et al*, 1996) include "flux adjustments" (also called "flux corrections") (Manabe and Stouffer, 1988; Sausen *et al*, 1988), which are prescribed additional fluxes of heat, water and momentum applied at the air-sea interface in order to maintain realistic sea-surface conditions in the control. As well as giving greater confidence in the simulation of climate change, a better control climatology offers an additional advantage regarding model evaluation against observations. If there are large errors in the surface climatology, there will probably be consequential errors in many other simulated quantities (precipitation, cloud, sea-ice, etc.); hence comparison with observations will not be so useful in helping us to assess the performance of these other aspects of the model.

The flux adjustments depend only on location and time of year, not on the model state, so are exactly the same in the control and perturbed climates. Thus they do not explicitly introduce any additional feedbacks. However, they do give cause for concern because they may locally have values which are large compared with the physically computed atmosphere-ocean fluxes. If the flux adjustment, which is independent of climate, constitutes an important part of the surface budget, there is a distortion of the relationship between state variables and fluxes. Also, there is an implied transport between areas of negative and positive flux adjustment, again independent of climate. Both of these may affect the climate response.

Some AOGCMs have been flux-adjusted in climate-change experiments (*e.g.* Cubasch *et al*, 1992; Manabe *et al*, 1992; Mitchell *et al*, 1995) and others not (*e.g.* Colman *et al*, 1995; Russell *et al*, 1995; Meehl and Washington, 1996). The climate response is qualitatively similar with and without, but there are differences in climate sensitivity and regional detail. It is not obvious whether any of these differences are due to flux adjustment, because the models differ in their formulations in many other respects which affect their climate response. To address this question, we have performed parallel climate-change experiments with and without flux adjustment using an AOGCM which is in all other respects exactly the same, allowing the first direct comparison of the response in the two cases. (A similar investigation has recently been carried out by Fanning and Weaver (1997) using an idealised AOGCM.) Here we report the basic results. Another paper will follow to describe the mechanisms in detail.

Model and experiments

We use a version of the Hadley Centre model known as HadCM2, which is normally run with heat and water flux adjustments to maintain realistic sea-surface temperatures and salinities, and has been used in various studies (Mitchell *et al*, 1995; Tett *et al*, 1996; Mitchell and Johns, 1997; Keen and Murphy, 1997). Johns *et al* (1997) describe the model, the spin-up, the derivation of the flux adjustments and the fluxadjusted control run, which we will refer to as "FC". FC has been run for over 1000 a and shows an upward drift in global-average surface air temperature of 0.16 mK a^{-1}. The global-average annual-average heat flux adjustment is 0.3 W m^{-2} (into the ocean), but locally it has a magnitude of up to 260 W m^{-2}.

An ensemble of four 80-year integrations were run with flux adjustment with CO_2 increasing at 1% per year compounded, reaching double its initial value after 70 years (Keen and Murphy, 1997). Radiative forcing increases linearly in time in these integra-

Figure 1: Timeseries of global-average temperature in the unfluxadjusted control run



tions, since it depends logarithmically on CO_2 concentration. The integrations began at different arbitrary points of FC, so developed differently in detail. The ensemble-average is the best estimate of the climate change, and the intraensemble variability gives information about the statistical uncertainty of the result. We refer to this anomaly ensemble as "FA".

An integration ("UC") without flux adjustment was begun from an arbitrary point of FC, and an ensemble ("UA") of four integrations, exactly like those of FA but without flux adjustment, were started from various points of UC. The earliest began after 30 years, to allow for an initial adjustment, which we describe next. Because there is a substantial drift in UC, the signal of climate change without flux adjustment was obtained by subtracting corresponding years of UC from the UA integrations. For FC the drift is negligible so a long-term mean can be used to characterise its climate.

Unfluxadjusted control climate

In the first decade of UC, the climate cools rapidly (Figure 1). Over years 11–15 the global-average surface air temperature is 2.2 K lower than in FC, and the sea-surface temperature (SST) 1.7 K lower. The cooling is not the same everywhere. It is fairly uniform between 45°N and 45°S, where the amounts of low and medium-level cloud increase substantially and high cloud moves downwards. High northern latitudes cool more strongly, connected with an expansion of the area of sea-ice and snow and an increase in global-average surface albedo. There is a warming in the Southern Ocean. After the initial phase, the temperature in UC settles into a long-term slow downward drift, being 2.6 K cooler in surface air temperature than FC after 100 years; this continues during the 360 years for

Figure 2: Timeseries of global-average temperature change in response to increasing CO_2 with and without flux adjustment



which UC has been run, and is fitted well by a linear trend of 1.9 mK a^{-1} for surface air temperature and 1.1 mK a^{-1} for SST. This drift is an order of magnitude larger than the warming tendency exhibited by FC.

It seems surprising that a global-average cooling as large as 2–3 K should result from a reduction in global-average heat input (the removed heat flux adjustment) of only 0.3 W m^{-2}, since a doubling of atmospheric CO_2, which produces a radiative forcing of ~ 3.5 W m^{-2} (after stratospheric adjustment), leads to a temperature rise of ~ 2.5 K (the IPCC "best estimate", Kattenberg *et al*, 1996). This comes about because the heat flux adjustment is *locally* large, so its removal causes large local temperature responses, but because there are geographically dependent feedbacks and long-range effects, there is not a uniform proportionality between local heat flux adjustments and temperature changes. Hence the positive and negative temperature changes do not balance to produce a small global average. The lack of proportionality can be seen in the small spatial correlations of the heat flux adjustment of about -0.2 with the surface air temperature change and -0.3 with the SST change. It appears that to eliminate the global-average temperature drift, we need the flux adjustments to be small *locally*, not just in the global average.

Climate change

The global-average temperature in both FA and UA rises fairly linearly with time, but less rapidly in UA (Figure 2). By the time when CO_2 has reached double its initial value (calculated as an average over years 61–80), the surface air temperature has risen by 1.76 ± 0.07 K in FA, but only 71% of this, 1.24 ± 0.06 K, in UA. (The uncertainty shown is the intraensemble standard deviation.) A comparison of zonal averages shows that the

Figure 3: Zonal-average global-average temperature change in response to increased CO_2 with and without flux adjustment



warming is reduced at all latitudes, by a fairly constant factor (Figure 3). The spatial correlation between the geographical distributions of surface air temperature change in FA and UA is 0.82, indicating the patterns are fairly similar. The correlation for SST is 0.69, which is smaller because part of the common signal in the surface air temperature is the land-sea contrast in the warming (greater over land). There are local differences, especially in northern high latitudes, which can partly be related to specific local differences in the climatologies of FC and UC.

When considering timescales of several years, the heat capacity of the atmosphere and mixed layer can be neglected. Therefore if Q is the imposed radiative forcing of the climate system,

$$Q = H + F \tag{1}$$

(see Figure 4). H is the increase in radiation to space at the top of the atmosphere (TOA), both longwave and shortwave, including feedbacks, resulting from the warming. F is the heat flux from the mixed layer into the deep ocean. When the climate regains a steady state, there must be no flux on average from the mixed layer to the deep ocean *i.e.* $F = 0 \Rightarrow Q = H$. Thus, H characterises the steady-state response of the climate system to Q, and F the transient behaviour. It may not be possible to attain a steady state in practice in an AOGCM, because the timescale on which the deep ocean equilibrates is millenia, and computing resources may not be available to perform such an integration.

Greenhouse gases inhibit the emission of longwave radiation to space, so the forcing Q they cause appears as a reduction in outgoing longwave radiation (OLR) at the TOA. Since H is an increase in TOA outgoing radiation, the change in the net downward flux at the TOA is $N = Q - H$. Assuming that H depends linearly on the temperature change ΔT of the climate system (in practice the global-average surface temperature), we write

Figure 4: Relationships between global-average fluxes



$H = \lambda\Delta T$, with λ a constant "climate response factor" (see *e.g.* Hansen *et al*, 1984; Dickinson, 1986). Hence $\lambda\Delta T = Q - N$. This enables us to obtain λ from N, Q and ΔT at any time; for the time of doubled CO_2 we find $\lambda = 1.32 \pm 0.06$ W m^{-2} K^{-1} for FA, 1.79 ± 0.10 W m^{-2} K^{-1} for UA (Table 1).

The conventional "climate sensitivity" $\Delta T_{2\times}$ is the global-average temperature difference between the control and a steady-state climate with a doubled concentration of CO_2. For a steady state $Q = H$, so if the forcing is $Q_{2\times}$, knowing λ we can obtain $\Delta T_{2\times}$ as $Q_{2\times}/\lambda$, without having integrated the model to a steady state. For FA, $\Delta T_{2\times} = 2.64 \pm 0.12$ K; for UA, 1.94 ± 0.19 K (Table 1). Both are within the generally accepted range (Kattenberg *et al*, 1996).

H is a combination of (Figure 4):

H_{LN} Change in clear-sky OLR (*i.e.* computed with no cloud) arising from the basic radiative response of the system to the higher temperature and from the water-vapour feedback.

H_{LC} Change in longwave cloud radiative forcing (CRF). Longwave CRF is the difference between clear-sky OLR and OLR with normal cloud, and is positive in the global average because clouds inhibit longwave radiation to space.

H_{SN} Change in clear-sky net downward shortwave (*i.e.* a reduction in upward shortwave), resulting from reduced surface albedo.

H_{SC} Change in shortwave cloud radiative forcing. Shortwave CRF is defined like longwave CRF, and is negative because clouds reflect sunlight.

Table 1: Comparison of climate response

	FA	UA
Change in global-average temperature ΔT/K	1.76	1.24
Radiative forcing $Q = Q_{2\times}$/W m^{-2}	3.47	3.47
Change in net radiation at TOA N/W m^{-2}	1.16	1.25
$H = Q - N = \lambda \Delta T$/W m^{-2}	2.31	2.22
λ/W m^{-2} K^{-1}	1.32	1.79
$\Delta T_{2\times}$/K	2.64	1.94
λ_{LN}/W m^{-2} K^{-1}	1.82	1.81
λ_{LC}/W m^{-2} K^{-1}	0.03	−0.28
λ_{SN}/W m^{-2} K^{-1}	0.71	0.85
λ_{SC}/W m^{-2} K^{-1}	−0.24	−0.55
$F = N = \kappa \Delta T$/W m^{-2}	1.16	1.25
κ/W m^{-2} K^{-1}	0.66	1.01

The first of these is defined upward, the others downward, so $H = H_{\mathrm{LN}} - H_{\mathrm{LC}} - H_{\mathrm{SN}} - H_{\mathrm{SC}}$. Each of them is assumed separately proportional to ΔT, so the combined $\lambda = \lambda_{\mathrm{LN}} - \lambda_{\mathrm{LC}} - \lambda_{\mathrm{SN}} - \lambda_{\mathrm{SC}}$. When $\lambda_{\mathrm{LC}}, \lambda_{\mathrm{SN}}, \lambda_{\mathrm{SC}}$ are positive, they give downward fluxes, which reinforce the radiative forcing, and constitute a positive feedback on the warming.

FA and UA have similar clear-sky longwave feedback λ_{LN} (Table 1), which means that the temperature difference between FC and UC is not large enough to have an important influence on the σT^4 term or the water-vapour feedback. The positive clear-sky shortwave feedback λ_{SN} is slightly stronger in UA than in FA, presumably because of the greater initial area of sea-ice and snow. The main difference in the response derives from the cloud feedbacks; the explanation must lie in the different cloud climatologies of FC and UC. λ_{SC} and λ_{LC} both become more negative, the most important influence being reduced positive feedbacks at low latitudes. For the shortwave, this could be because UC has much more low and medium-level cloud than FC. Consequently, the reduction of high cloud resulting from the warming has less effect on the planetary albedo. For the longwave, a possible explanation is that the increase in the upper-level cloud water path is less in UA because the temperature is lower.

Assuming the heat capacity of the mixed layer and atmosphere is small, the net downward flux N at the TOA is nearly equal to the net heat flux F into the deep ocean (Figure 4). Timeseries of N show that it grows roughly linearly in time (Figure 5). Since ΔT also increases approximately linearly in time, there is a proportionality $F = \kappa \Delta T$. We do not expect this *a priori*, and it does not have the general applicability of $H = \lambda \Delta T$. It obviously fails as the system approaches a steady state, when $F \to 0$ but ΔT tends to a non-zero final value. Nonetheless, it is useful in the present analysis.

It is surprising that F is very similar in FA and UA; again, there is no reason to expect this, which means that the heat taken up by the deep ocean is the about same in the two experiments, despite the fact that the surface warming is 30% less in UA. In fact, the increase in heat content of the entire ocean averaged over years 61–80 in UA is only 8% less than in FA. A calculation of the distribution of this heat with depth shows that, on average, it penetrates more deeply in UA, meaning that the vertical mixing processes are more effective in the unfluxadjusted ocean (Figure 6). The mechanisms for this will be discussed in a later paper.

Figure 5: Timeseries of global-average net downward radiation at the top of the atmosphere



Figure 6: Storage of additional heat in the ocean as a result of increased CO_2 with and without flux adjustment



The constant κ quantifies the transfer of heat from the surface to the deep ocean. Since F is the same but ΔT is less in UA, κ must be greater, consistent with the conclusion drawn from the vertical distribution of heat uptake. Calculating $\kappa = F/\Delta T$ yields 0.66 ± 0.04 W m^{-2} K^{-1} in FA, 1.01 ± 0.16 W m^{-2} K^{-1} in UA.

In terms of ΔT, Equation (1) becomes

$$Q = (\lambda + \kappa)\Delta T. \tag{2}$$

This shows that the surface temperature is determined by the effectiveness of heat loss to space and into the deep ocean. Because λ and κ are bigger in the unfluxadjusted model, ΔT is smaller. The differences between the two models in λ and κ are similar, so the smaller climate sensitivity and the increased vertical transport are of comparable importance in reducing ΔT.

Conclusion

Although the global-average flux adjustment is small, substantial local and global-average climate drift occurs in HadCM2 without it. The drift is amplified by local feedbacks, suggesting that the flux adjustment has to be small locally, as well as on average, to eliminate global-average drift. From the comparison of response experiments, we find that global-average temperature change resulting from increasing CO_2 is less in our model without flux adjustment than with, everything else in the model being the same. The geographical patterns of temperature change are generally similar. The altered response is not, however, due directly to the flux adjustment, but to the considerably different initial states of the atmosphere and ocean; it comes about mainly through a reduced climate sensitivity caused by differences in the cloud climatology, and increased effectiveness of vertical heat transport in the ocean. Insofar as the fluxadjusted initial state is closer to observations, this comparison provides support for the use of the technique of flux adjustment; we have more confidence in the climate response from the fluxadjusted model because its control state is more realistic. At this stage in the analysis we cannot rule out the possibility that the flux adjustment does also directly influence the climate response, although apparently it does not introduce any gross distortion. Our best strategy remains to try to improve the model to reduce the flux adjustments, repeating the comparison of climate response with and without the technique until they converge.

Acknowledgements

This work was supported by the UK Department of the Environment under contract PECD 7/12/37.

References

Colman, R. A., S. B. Power, B. J. McAvaney, and R. R. Dahni, 1995: A non-flux-corrected transient CO_2 experiment using the BMRC coupled atmosphere/ocean GCM. *Geophys. Res. Lett.*, **22**, 3047–3050.

Cubasch, U., K. Hasselmann, H. Höck, E. Maier-Reimer, U. Mikolajewicz, B. D. Santer, and R. Sausen, 1992: Time-dependent greenhouse warming computations with a coupled ocean-atmosphere model. *Climate Dynamics*, **8**, 55–69.

Dickinson, R. E., 1986: How will climate change? *The Greenhouse Effect, Climate Change and Ecosystems*, Bolin, B., B. R. Doos, J. Jaeger, and R. A. Warrick, Eds., SCOPE Report 29, John Wiley and Sons, 206–270.

Fanning, A. F., and A. J. Weaver, 1997. On the role of flux adjustments in an idealized coupled climate model. Submitted to *Clim. Dyn.*

Gates, W. L., A. Henderson-Sellers, G. J. Boer, C. K. Folland, A. Kitoh, B. J. McAvaney, F. Semazzi, N. Smith, A. J. Weaver, and Q.-C. Zeng, 1996: Climate models—Evaluation. *Climate Change 1995. The Science of Climate Change*, Houghton, J. T., L. G. Meira Filho, B. A. Callandar, N. Harris, A. Kattenberg, and K. Maskell, Eds., Cambridge University Press, 235–284.

Hansen, J. E., A. Lacis, D. Rind, L. Russell, et al., 1984: Climate sensitivity analysis of feedback mechanisms. *Climate Processes and Climate Sensitivity*, Hansen, J. E., and T. Takahashi, Eds., Geophysical Monograph 29, American Geophysical Union, 130–163.

Johns, T. C., R. E. Carnell, J. F. Crossley, J. M. Gregory, J. F. B. Mitchell, C. A. Senior, S. F. B. Tett, and R. A. Wood, 1997. The second Hadley Centre coupled ocean-atmosphere GCM: Model description, spinup and validation. To appear in *Clim. Dyn.*

Kattenberg, A., F. Giorgi, H. Grassl, G. A. Meehl, J. F. B. Mitchell, R. J. Stouffer, T. Tokioka, A. J. Weaver, and T. M. L. Wigley, 1996: Climate models—projections of future climate. *Climate Change 1995. The Science of Climate Change*, Houghton, J. T., L. G. Meira Filho, B. A. Callander, N. Harris, A. Kattenberg, and K. Maskell, Eds., Cambridge University Press, 285–358.

Keen, A. B., and J. M. Murphy, 1997. Influence of natural variability and the cold start problem on the simulated transient response to increasing CO_2. To be submitted to *Clim. Dyn.*

Manabe, S., and R. J. Stouffer, 1988: Two stable equilibria of a coupled ocean-atmosphere model. *J. Climate*, **1**, 841–866.

——, M. J. Spelman, and R. J. Stouffer, 1992: Transient response of a coupled ocean-atmosphere model to gradual changes of atmospheric CO_2. Part II. seasonal response. *J. Climate*, **5**, 105–126.

Meehl, G. A., and W. M. Washington, 1996: El Niño-like climate change in a model with increased atmospheric CO_2 concentrations. *Nature*, **382**, 56–60.

Mitchell, J. F. B., and T. C. Johns, 1997: On the modification of global warming by sulphate aerosols. *J. Climate*, **10**, 245–267.

——, ——, J. M. Gregory, and S. F. B. Tett, 1995: Climate response to increasing levels of greenhouse gases and sulphate aerosols. *Nature*, **376**, 501–504.

Russell, G. L., J. R. Miller, and D. Rind, 1995: A coupled atmosphere-ocean model for transient climate change studies. *Atmos. Ocean*, **33**, 683–730.

Sausen, R., K. Barthel, and K. Hasselmann, 1988: Coupled ocean-atmosphere models with flux corrections. *Clim. Dyn.*, **2**, 154–163.

Tett, S. F. B., T. C. Johns, and J. F. B. Mitchell, 1996. Global and regional variability in a coupled AOGCM. Submitted to *Clim. Dyn.*

CLIMATE RESEARCH TECHNICAL NOTES

CRTN 1	Oct 1990	Estimates of the sensitivity of climate to vegetation changes using the Penman-Monteith equation. P R Rowntree
CRTN 2	Oct 1990	An ocean general circulation model of the Indian Ocean for hindcasting studies. D J Carrington
CRTN 3	Oct 1990	Simulation of the tropical diurnal cycle in a climate model. D P Rowell
CRTN 4	Oct 1990	Low frequency variability of the oceans. C K Folland, A Colman, D E Parker and A Bevan
CRTN 5	Dec 1990	A comparison of 11-level General Circulation Model Simulations with observations in the East Sahel. K Maskell
CRTN 6	Dec 1990	Climate Change Prediction. J F B Mitchell and Qing-cun Zeng
CRTN 7	Jan 1991	Deforestation of Amazonia - modelling the effects of albedo change. M F Mylne and P R Rowntree
CRTN 8	Jan 1991	The role of observations in climate prediction and research. D J Carson
CRTN 9	Mar 1991	The greenhouse effect and its likely consequences for climate change. D J Carson
CRTN 10	Apr 1991	Use of wind stresses from operational N.W.P. models to force an O.G.C.M. of the Indian Ocean. D J Carrington
CRTN 11	Jun 1991	A new daily Central England Temperature series, 1772-1991. D E Parker, T P Legg and C K Folland
CRTN 12	Jul 1991	Causes and predictability of Sahel rainfall variability. D P Rowell, C K Folland, K Maskell, J A Owen, M N Ward

CLIMATE RESEARCH TECHNICAL NOTES

CRTN 13	Jul 1991	Modelling changes in climate due to enhanced CO2, the role of atmospheric dynamics, cloud and moisture. C A Senior, J F B Mitchell, H Le Treut and Z-X Li
CRTN 14	Sep 1991	Sea temperature bucket models used to correct historical SST data in the Meteorological Office. C K Folland
CRTN 15	Aug 1991	Modelling climate change, and some potential effects on agriculture in the U.K. P R Rowntree, B A Callander and J Cochrane
CRTN 16	Aug 1991	The Boreal Forests and Climate G Thomas and P R Rowntree
CRTN 17	Aug 1991	Development of a Stratosphere-Troposphere Data Assimilation System. R Swinbank.
CRTN 18	Sep 1991	A study of asynchronous coupling using a simple climate model. M K Davey.
CRTN 19	Sep 1991	The Oceanic Carbon Cycle. N K Taylor
CRTN 20	Nov 1991	Worldwide ocean-atmosphere surface fields in Sahel wet and dry years using provisionally corrected surface wind data. M N Ward
CRTN 21	Dec 1991	Coupled tropical ocean global atmosphere models at the UKMO. M K Davey, C Gordon, S Ineson and S Lawrence
CRTN 22	Dec 1991	Empirical parameterisation of tropical ocean-atmosphere coupling. M Allen, M K Davey, D L T Anderson and P D Killworth
CRTN 23	Feb 1992	The temporal evolution of Equatorial Currents in the Indian Ocean D L T Anderson, D J Carrington, R A Corry and C Gordon
CRTN 24	Feb 1992	Stratospheric analyses provided by the U.K. Meteorological Office M J Bailey, A O'Neill and V D Pope

CLIMATE RESEARCH TECHNICAL NOTES

CRTN 25	Feb 1992	Modelling interannual variability in the Indian Ocean using momentum fluxes from the UKMO and ECMWF operational weather analyses. D L T Anderson and D J Carrington
CRTN 26	Mar 1992	A GCM simulation of the impact of Amazonian deforestation on climate using an improved canopy representation J Lean and P R Rowntree
CRTN 27	Mar 1992	The parameterization of rainfall interception in GCMs A J Dolman and D Gregory
CRTN 28	Jun 1992	Development of worldwide marine data eigenvectors since 1985 A W Colman
CRTN 29	Jun 1992	A tropical ocean model with reduced physics M K Davey
CRTN 30	Jun 1992	International Temperature Workshop, Boulder, Colorado, USA, 16 January 1992 Edited by D E Parker
CRTN 31	Jul 1992	Simulation of clear-sky outgoing longwave radiation over the oceans using operational analyses A Slingo and M J Webb
CRTN 32	Sep 1992	A prediction of the transient response of climate J M Murphy
CRTN 33	Nov 1992	LEPS scores for assessing climate model simulations and long-range forecasts. C K Folland
CRTN 34	Jan 1993	Stratospheric Data Assimilation System Guide Editer: R Swinbank
CRTN 35	Mar 1993	A Stratosphere-Troposphere Data Assimilation System R Swinbank and A O'Neill
CRTN 36	Mar 1993	Validation of hydrological schemes for climate models against catchment data. P R Rowntree and J Lean

CLIMATE RESEARCH TECHNICAL NOTES

CRTN 37	Apr 1993	Modelling of palaeoclimates: Examples from the recent past J F B Mitchell
CRTN 38	May 1993	A simulation of seasonality in ENSO forecast skill M K Davey, D L T Anderson and S Lawrence
CRTN 39	Jul 1993	ENSO Prediction experiments using a simple ocean-atmosphere model D-H Wu, D L T Anderson and M K Davey
CRTN 40	Sep 1993	Diagnosis of dynamic sea-surface and sea-level changes from the Cox ocean model J M Gregory
CRTN 41	Sep 1993	Seasonal variations of the clear-sky greenhouse effect: the role of changes in atmospheric temperatures and humidities M J Webb, A Slingo and G L Stephens
CRTN 42	Nov 1993	Sea-level changes under increasing atmospheric CO_2 in a transient coupled ocean-atmosphere GCM experiment. J M Gregory
CRTN 43	Dec 1993	Global and regional patterns of climate change: recent predictions for the Arctic P R Rowntree
CRTN 44	Feb 1994	The effect of changing horizontal diffusion in the atmospheric version of the unified climate model C D Hall and R A Stratton
CRTN 45	Mar 1994	Simulation of El-Niño/Southern Oscillation like variability in a global AOGCM and its response to CO_2 increase. S F B Tett
CRTN 46	Apr 1994	Global data required for monitoring climate change. D E Parker and C K Folland
CRTN 47	May 1994	Seasonal uptake of anthropogenic CO_2 in an ocean general circulation model. N K Taylor

CLIMATE RESEARCH TECHNICAL NOTES

CRTN 48	Jun 1994	A tropic-wide oscillation of boreal summer rainfall and patterns of sea-surface temperature. M N Ward, K Maskell, C K Folland, D P Rowell and R Washington
CRTN 49	Jun 1994	Simulation of the tropical Pacific using a simplified ocean model M A Balmaseda, D L T Anderson and M K Davey
CRTN 50	Sep 1994	Simulation of global mean temperature using a box-diffusion climate model P R Rowntree
CRTN 51	Sep 1994	Seasonal dependence of ENSO prediction skill M A Balmaseda, M K Davey and D L T Anderson
CRTN 52	Nov 1994	Simulation of the Indian Monsoon and Tropical Intraseasonal Variability by a General Circulation Model P M Inness and D Gregory
CRTN 53	Mar 1995	A comparison of modelled surface fluxes with climatological estimates. C Gordon and D K Wright
CRTN 54	Mar 1995	The representation of moist convection in atmospheric models. D Gregory
CRTN 55	Mar 1995	A consistent treatment of the evaporation of rain and snow for use in large-scale models D Gregory
CRTN 56	Apr 1995	Workshop on simulations of the Climate of the Twentieth Century using GISST, 28-30 November 1994, Hadley Centre, Bracknell, UK Edited by C K Folland and D P Rowell
CRTN 57	May 1995	Enhanced shortwave cloud radiative forcing due to anthropogenic aerosols. S E Swartz and A Slingo
CRTN 58	May 1995	The simulation of the tropical oceans in models of different horizontal resolution. C Gordon, D K Wright and C M Roberts
CRTN 59	Sep 1995	The water budget of middle latitude continental regions - a modelling and observational study P R Rowntree
CRTN 60	Sep 1995	Intraseasonal variability of the Indian summer monsoon simulated by the Hadley Centre climate model K Ashok, D Gregory and P M Inness

CLIMATE RESEARCH TECHNICAL NOTES

CRTN 61	Sep 1995	Climate simulations with the Unified Model: AMIP runs C D Hall, R A Stratton and M L Gallani
CRTN 62	Nov 1995	Validation of surface parameters over the oceans in climate simulations with the unified model. C D Hall
CRTN 63	Dec 1995	The GISST2.2 sea surface temperature and sea-ice climatology D E Parker, M Jackson and E B Horton
CRTN 64	Dec 1995	Parametrization of momentum transport by convection. II: Tests in single column and general circulation models. D Gregory, R Kershaw and P M Inness
CRTN 65	Jan 1996	Understanding the sensitivity of a GCM simulation of Amazonian deforestation to the specification of vegetation and soil characteristics. J Lean and P R Rowntree
CRTN 66	Jan 1996	On the efficient calculation of infra-red fluxes and cooling rates using the two-stream equations. J M Edwards
CRTN 67	Jan 1996	The second Hadley Centre coupled ocean-atmosphere GCM: Model description, spinup and validation. TC Johns,RE Carnell,JF Crossley,JM Gregory, JFB Mitchell,CA Senior,SFB Tett and RA Wood
CRTN 68	Feb 1996	The sensitivity of climate simulations to the specification of mixed phase clouds D Gregory and D Morris
CRTN 69	Apr 1996	Using an ensemble of Multi-Decadal GCM simulations ro assess potential seasonal predictability D P Rowell
CRTN 70	May 1996	North Atlantic and European Seasonal Predictability using an Ensemble of Multi-Decadal AGCM Simulations J R Davies, D P Rowell and C K Folland
CRTN 71	Jun 1996	A description of the Second Hadley Centre Coupled Model (HADCM2) T C Johns
CRTN 72	Jul 1996	Palaeoclimate Modelling Intercomparison: UKMO GCM simulations for 6kBP and 21kBP C D Hewitt and J F B Mitchell

CLIMATE RESEARCH TECHNICAL NOTES

CRTN 73	Aug 1996	ENSO responses and low frequency weather variability in the North Pacific/American sector 1949-93 A C Renshaw, D P Rowell and C K Folland
CRTN 74	Sep 1996	Version 2.2 of the Global sea-Ice and Sea Surface Temperature data set, 1903-1994 N A Rayner, E B Horton, D E Parker, C K Folland and R B Hackett
CRTN 75	Sep 1996	A new gravity wave drag scheme incorporating anisotropic orography and low level wave breaking : Impact upon the climate of the UK Met Office Unified Model D Gregory, G J Shutts and J R Mitchell
CRTN 76	Nov 1996	Middle atmosphere variability in the UKMO Unified Model R Swinbank, W Lahoz, C Douglas, A Heaps, R Brugge, W Norton, A O'Neill, D Podd
CRTN 77	Dec 1996	A high resolution AMIP run using the Hadley Centre model HadAM2b R A Stratton
CRTN 78	Jan 1997	The climate response to CO_2 of the Hadley Centre coupled AOGCM with and without flux adjustment J M Gregory and J F B Mitchell